

<b>National Imaging Associates, Inc.*</b>	
<b>Clinical Guidelines</b> <b>STRESS ECHOCARDIOGRAPHY</b>	<b>Original Date: February 2010</b>
<b>CPT Codes: 93350, 93351, +93320, +93321, +93325, +93352, +93356</b>	<b>Last Revised Date: <del>March-February</del> 202<del>2</del><sup>1</sup></b>
<b>Guideline Number: NIA_CG_026</b>	<b>Implementation Date: January 202<del>3</del><sup>2</sup></b>

## GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

[This guideline is for stress imaging, specifically Stress Echocardiography \(SE\) with appropriate preference for suitable alternatives, such as an exercise treadmill exam without imaging, when more suitable, unless otherwise stated \(refer to Overview section\).](#)

## INDICATIONS for STRESS ECHO

### SUSPECTED CORONARY ARTERY DISEASE (CAD)

#### Symptomatic patients without known CAD (use [Diamond Forrester table](#))

- Low or intermediate pretest probability, and electrocardiogram (ECG) is uninterpretable
- High pretest probability
- Repeat testing in patient with new or worsening symptoms and negative result at least one year ago AND meets one of the criteria above

#### Asymptomatic patients without known CAD

- Previously unevaluated ECG evidence of possible myocardial ischemia including ischemic ST segment or T wave abnormalities ([See Overview section](#))
- Previously unevaluated pathologic Q waves ([See Overview Section](#))
- Previously unevaluated complete left bundle branch block

\* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

- ~~Those with a calcium score >400, not previously evaluated History of diabetes mellitus, >40 years old, with calcium score >400<sup>±</sup> (add cp 2021 reference) (Budoff, 2016)~~

### **ABNORMAL CALCIUM SCORES (CAC)<sup>1-5</sup>**

- ASYMPTOMATIC patient with a calcium score >400, not previously evaluated
- SYMPTOMATIC patient with prior CAC ≥100

### **INCONCLUSIVE CAD EVALUATION WITHIN THE PAST 2 YEARS AND OBSTRUCTIVE CAD REMAINS A CONCERN**

- Exercise stress ECG with low risk Duke treadmill score ≥5, but patient's current symptoms indicate an intermediate or high pretest probability
- Exercise stress ECG with an intermediate Duke treadmill score
- Intermediate coronary computed tomography angiography (CCTA) defined as:
  - ~~430~~ -70% lesion

### **FOLLOW-UP OF PATIENTS POST CORONARY REVASCULARIZATION (PCI or CABG)<sup>6</sup>** **(Doherty, 2019)**

- **Asymptomatic, follow-up stress imaging** (MPI or SE), at a minimum of 2 years post coronary artery bypass grafting (CABG), or percutaneous coronary intervention (PCI), whichever is later, is appropriate for patients with a history of silent ischemia or a history of a prior left main stent<sup>3</sup> ~~(Wolk, 2014)~~
- OR
- For patients with high occupational risk including any of the following:
  - Associated with public safety
  - Airline and boat pilots
  - Bus and train drivers
  - Bridge and tunnel workers/toll collectors
  - Police officers and firefighters
- **New, recurrent, or worsening symptoms post coronary revascularization** is an indication for stress imaging

### **FOLLOW-UP OF KNOWN CAD**

~~Symptomatic Low or intermediate pretest probability chest pain with coronary disease defined by a Coronary calcium score ≥100 is ok for stress imaging~~

- **Routine follow-up of asymptomatic or stable symptoms** when last invasive or non-invasive assessment of coronary disease showed hemodynamically significant CAD (ischemia on stress test or FFR ≤ 0.80 or stenosis greater than or equal to 70% of a major vessel), over two years ago without intervening coronary revascularization, is an appropriate indication for stress imaging (MPI or SE)

## SPECIAL DIAGNOSTIC CONDITIONS REQUIRING CORONARY EVALUATION

- Prior acute coronary syndrome (with documentation in MD notes), within last three months, without a prior stress test or coronary angiography performed since that time
- Newly diagnosed systolic heart failure or diastolic heart failure, with reasonable suspicion of cardiac ischemia (prior events, risk factors), with symptoms or signs of ischemia unless invasive coronary angiography is immediately planned<sup>1, 7</sup> (ref) 1, 6, 7 of MPI guideline and new reference needs adding at end (AUC rec) Newly diagnosed systolic heart failure (EF < 50%), when invasive coronary angiography has not been performed, especially when symptoms or signs of ischemia are present or suspected such as:
  - Chest pain
  - EKG changes such as new ST segment depression or T wave inversions
  - New wall motion abnormalities
- Ventricular arrhythmias:
  - Sustained ventricular tachycardia (VT) > 100 bpm, ventricular fibrillation (VF), or exercise-induced VT, when invasive coronary arteriography has not been performed<sup>8</sup>
  - Nonsustained VT, multiple episodes, each ≥ 3 beats at ≥ 100 bpm, frequent ~~VPC's~~ PVCs (defined as greater than or equal to 30/hour on remote monitoring), when an exercise ECG cannot be performed<sup>9</sup> (Zimetbaum, 2018)
- For intermediate and high risk global patients who require ~~Prior to~~ initiation of Class IC antiarrhythmic drugs. It can be performed annually thereafter until discontinuation of drug use ~~initiation (Propafenone or Flecainide), as well as annually in intermediate and high global risk patients~~<sup>10</sup>
- Hemodynamic assessment of ischemia in one of the following documented conditions:
  - Anomalous coronary arteries in an asymptomatic individual without prior stress echocardiography<sup>11</sup>;
  - Myocardial bridging of a coronary artery (perform with exercise stress)<sup>12</sup>;
- Coronary aneurysms in Kawasaki's disease<sup>13</sup>
- Following radiation therapy to the anterior or left chest, at 5 years post initiation and every 5 years thereafter<sup>14</sup>

## CHRONIC VALVULAR DISEASE

### Evaluation with Inclusion of Doppler<sup>15-18</sup>

(Baumgartner, 2017; Bonow, 2020; Nishimura, 2014; Steiner, 2017)

- Dobutamine SE for the evaluation of aortic stenosis and flow (contractile) reserve in symptomatic patients with severe aortic stenosis by calculated valve area, low flow / low gradient, and ejection fraction < 50%
- Exercise echo Doppler evaluation for mitral stenosis (MS) if there is:

- Exertional shortness of breath which suggests the amount of MS is worse than is seen on the resting echocardiogram
- Exercise echo Doppler evaluation for mitral regurgitation (MR) if there is:
  - Exertional shortness of breath which suggests the amount of MR is worse than is seen on the resting echocardiogram; **OR**
  - The echocardiogram is not able to distinguish whether the MR is moderate or severe in a patient that is asymptomatic
- For symptomatic patients with HCM, who do not have resting or provokable outflow tract gradient  $\geq 50$  mm Hg on TTE, for detection and quantification of dynamic LVOT obstruction<sup>19</sup> ~~(Ommen, 2020)~~
- For asymptomatic patients with HCM who do not have a resting or provokable outflow tract gradient  $\geq 50$  mm Hg on TTE (Class 2A)

#### **PRIOR TO ELECTIVE NON-CARDIAC SURGERY<sup>20-23</sup>**

~~(Fleischer, 2014; Patel, 2015)~~

- An ~~i~~ntermediate or high risk surgery with of one or more risk factors (see below), AND documentation of an inability to walk (or  $<4$  METs) AND there has not been an imaging stress test **within 1 year**<sup>22, 24, 25</sup> ~~within 1 year\*~~ (see reference below)
  - Risk factors: history of ischemic heart disease, history of congestive heart failure, history of cerebrovascular disease, preoperative treatment with insulin, and preoperative serum creatinine  $>2.0$  mg/dL.
  - Surgical Risks:
    - High risk surgery: Aortic and other major vascular surgery, ~~p~~Peripheral vascular surgery, ~~a~~Anticipated prolonged surgical procedures associated with large fluid shifts and/or blood loss
    - Intermediate risk surgery: Carotid endarterectomy, ~~h~~Head and neck surgery, ~~i~~ntraperitoneal and intrathoracic surgery, ~~o~~Orthopedic surgery, ~~p~~Prostate surgery
    - Low risk surgery: Endoscopic procedures, superficial procedure, cataract surgery, ~~b~~Breast surgery

~~Patients who would otherwise not be planned for a non-invasive coronary evaluation, but are referred for preoperative cardiac evaluation, are eligible for SE if **ALL 4** criteria are met:~~

~~Surgery is supra-inguinal vascular, intrathoracic, or intra-abdominal; **AND**~~

~~The patient has **at least one** of these additional cardiac complication risk factors:~~

~~Ischemic Heart Disease~~

~~History of stroke or transient ischemic attack (TIA)~~

~~History of congestive heart failure (CHF) or ejection fraction  $\leq 35\%$~~

~~Insulin-requiring diabetes mellitus~~

~~Creatinine  $\geq 2.0$  mg/dl~~

~~**AND**~~

~~The patient has limited functional capacity ( $<4$  metabolic equivalents) such as one of the following (would likely be requested as MPD):~~

~~Cannot take care of their ADLs which include but not limited to:  
Independently eating, bathing or ambulating  
Cannot walk 2 blocks on level ground  
Cannot climb 1 flight of stairs~~

~~AND~~

~~There has not been a conclusive stress evaluation, CTA, or heart catheterization within the past year, and the results would be likely to preclude proceeding with the intended surgery~~

- Planning for any organ or stem cell transplantation is an indication for preoperative stress imaging MPI, if there has not been a conclusive stress evaluation, CTA, or heart catheterization within the past year, at the discretion of the transplant service.<sup>23, 26</sup>

~~Planning for solid organ transplantation (liver or kidney), is an indication for preoperative dobutamine SE, if there has not been a conclusive stress evaluation, CTA, or heart catheterization within the past year and with  $\geq 3$  of the following risk factors<sup>26</sup> (Lentine, 2012):~~

~~Age  $> 60$~~

~~Smoking~~

~~Hypertension~~

~~Dyslipidemia~~

~~Left ventricular hypertrophy~~

~~$> 1$  year on dialysis (for renal transplant patients)~~

~~Diabetes mellitus~~

~~Prior ischemic heart disease~~

## POST CARDIAC TRANSPLANTATION

- Annually, for the first five years post cardiac transplantation, in a patient not undergoing invasive coronary arteriography
- After the first five years post cardiac transplantation, patients with documented transplant coronary vasculopathy can be screened annually unless invasive coronary arteriography is planned
- ~~Annually, for the first five years post cardiac transplantation, in a patient not undergoing invasive coronary arteriography~~
- ~~After the first five years post cardiac transplantation, patients with transplant coronary vasculopathy can be screened annually with **ONE** of the following:~~
  - ~~○ MPI~~
  - ~~○ SE~~
  - ~~○ Left heart catheterization~~

---

## BACKGROUND

Stress echocardiography (SE) refers to ultrasound imaging of the heart during exercise electrocardiography (ECG) testing, during which visualized wall motion abnormalities can

provide evidence of potential significant coronary artery disease (CAD).

While drug-induced stress with dobutamine can be an alternative to exercise stress testing in patients who are unable to exercise, this guideline does not require use of this modality. Hence, reference in this document to SE predominantly refers to exercise stress echocardiography.

Although SE provides comparable accuracy without radiation risk, relative to myocardial perfusion imaging (MPI), scenarios which do not permit effective use of SE might be better suited for stress imaging with MPI, cardiovascular magnetic resonance imaging (CMR) or positron emission tomography (PET), or coronary computed tomography angiography (CCTA).

**Stable patients without known CAD** fall into 2 categories<sup>3, 23, 27</sup>:

~~(Fihn, 2012; Montalescot, 2013; Wolk, 2014)~~

- **Asymptomatic patients**, for whom Global Risk of CAD events can be determined from coronary risk factors using calculators available online (see Websites for [Global Cardiovascular Risk Calculators](#) section)
- **Symptomatic patients**, for whom we estimate the Pretest Probability that their chest-related symptoms are due to clinically significant CAD (see below):

#### **The 3 Types of Chest Pain or Discomfort:**

- **Typical Angina (Definite)** is defined as including all **3** of these characteristics:
  - Substernal chest pain or discomfort with characteristic quality and duration, such as:
    - Pressure-like
    - Radiating
    - Dull or aching
  - Provoked by exertion or emotional stress
  - Relieved by rest and/or nitroglycerine
- **Atypical Angina (Probable)** has only **2** of the above characteristics
- **Nonanginal Chest Pain/Discomfort** has only **0-1** of the above characteristics

The medical record should provide enough detail to establish the type of chest pain. From those details, ~~t~~The Pretest Probability of obstructive CAD is estimated from the Diamond Forrester Table ~~Once the type of chest pain has been established from the medical record, the Pretest Probability of obstructive CAD is estimated from the Diamond Forrester Table~~ below, recognizing that in some cases multiple additional coronary risk factors could increase pretest probability<sup>3, 27</sup> ~~(Fihn, 2012; Wolk, 2014)~~:

## Diamond Forrester Table

Age (Years)	Gender	Typical/Definite Angina Pectoris	Atypical/Probable Angina Pectoris	Nonanginal Chest Pain
≤ 39	Men	Intermediate	Intermediate	Low
	Women	Intermediate	Very low	Very low
40 – 49	Men	High	Intermediate	Intermediate
	Women	Intermediate	Low	Very low
50 – 59	Men	High	Intermediate	Intermediate
	Women	Intermediate	Intermediate	Low
≥ 60	Men	High	Intermediate	Intermediate
	Women	High	Intermediate	Intermediate

- **Very low:** < 5% pretest probability of CAD, usually not requiring stress evaluation<sup>27</sup> (Fihn, 2012)
- **Low:** 5 - 10% pretest probability of CAD
- **Intermediate:** 10% - 90% pretest probability of CAD
- **High:** > 90% pretest probability of CAD

## OVERVIEW

MPI may be performed without diversion to SE in any of the following<sup>3, 28</sup> (Henzlova, 2016; Wolk, 2014):

- Inability to exercise
  - Physical limitations precluding ability to exercise for at least 3 full minutes of Bruce protocol
  - BMI > 40kg/m<sup>2</sup>
  - Limited functional capacity (< 4 metabolic equivalents) **such as one** of the following:
    - Cannot take care of their activities of daily living (ADLs) or ambulate
    - Cannot walk 2 blocks on level ground
    - Cannot climb 1 flight of stairs
    - Cannot vacuum, dust, do dishes, sweep, or carry a small grocery bag
- Other Comorbidities
  - Severe chronic obstructive pulmonary disease with pulmonary function test (PFT) documentation, severe shortness of breath on minimal exertion, or requirement of home oxygen during the day
  - Poorly controlled hypertension, with systolic BP > 180 or Diastolic BP > 120 (and clinical urgency not to delay MPI)
- ECG and Echo-Related Baseline Findings
  - Prior cardiac surgery (coronary artery bypass graft or valvular)
  - Documented poor acoustic imaging window

- Left ventricular ejection fraction  $\leq 40\%$
- Pacemaker or ICD
- Persistent atrial fibrillation
- Resting wall motion abnormalities that would make SE interpretation difficult
- Complete LBBB

- Risk-related scenarios

- High pretest probability in suspected CAD
- Intermediate or high global risk in patients requiring type IC antiarrhythmic drugs (prior to initiation of therapy and annually)
- Arrhythmia risk with exercise

- Previously unevaluated pathologic Q waves (in two contiguous leads) waves defined as the following:

- > 40 ms (1 mm) wide
- > 2 mm deep
- > 25% of depth of QRS complex

#### Risk-related scenarios

High pretest probability in suspected CAD

Intermediate or high global risk in patients requiring type IC antiarrhythmic drugs (prior to initiation of therapy and annually)

Arrhythmia risk with exercise

## ECG Stress Test Alone versus Stress Testing with Imaging

Prominent scenarios suitable for an ECG stress test WITHOUT imaging (i.e., exercise treadmill ECG test) are inferred from the guidelines presented above, often requiring that the patient can exercise for at least 3 minutes of Bruce protocol with achievement of near maximal heart rate AND has an interpretable ECG for ischemia during exercise<sup>3</sup> ~~(Wolk, 2014)~~:

- The (symptomatic) low or intermediate pretest probability patient who is able to exercise and has an interpretable ECG
- ~~The (asymptomatic) high global risk patient who can exercise and has an interpretable ECG~~
- The patient who is under evaluation for exercise-induced arrhythmia<sup>8</sup> ~~(Al-Khatib, 2017)~~
- For the evaluation of syncope or presyncope during exertion<sup>29</sup>
- The patient who requires an entrance stress test ECG for a cardiac rehab program or for an exercise prescription.

## Duke Exercise ECG Treadmill Score<sup>30</sup> ~~(Mark, 1987)~~

Calculates risk from ECG treadmill alone:



- The equation for calculating the Duke treadmill score (DTS) is:  $DTS = \text{exercise time in minutes} - (5 \times \text{ST deviation in mm or 0.1 mV increments}) - (4 \times \text{exercise angina score})$ , with angina score being 0 = none, 1 = non-limiting, and 2 = exercise-limiting.
- The score typically ranges from - 25 to + 15. These values correspond to low-risk (with a score of  $\geq + 5$ ), intermediate risk (with scores ranging from - 10 to + 4), and high-risk (with a score of  $\leq -11$ ) categories.

An uninterpretable baseline ECG includes<sup>27</sup> ~~(Fihn, 2012)~~:

- ST segment depression 1 mm or more; (not for non-specific ST- T wave changes)
- Ischemic looking T wave -- at least 2.5 mm inversions (excluding V1 and V2)
- LVH, pre excitation pattern such as WPW, a ventricular paced rhythm, or left bundle branch block
- Digitalis use
- Resting HR under 50 bpm on a medication, such as beta-blockers or calcium channel blockers, that is required for patient's treatment and cannot be stopped, with an anticipated suboptimal workload

## Global Risk of Cardiovascular Disease

**Global risk** of CAD is defined as the probability of manifesting cardiovascular disease over the next 10 years and refers to **asymptomatic** patients without known cardiovascular disease. It should be determined using one of the risk calculators below. A high risk is considered greater than a 20% risk of a cardiovascular event over the ensuing 10 years. High global risk by itself generally lacks scientific support as an indication for stress imaging. There are rare exemptions, such as patients requiring IC antiarrhythmic drugs, who might require coronary risk stratification prior to initiation of the drug. ~~or patients with a CAC score > 400 Agatston units, when global risk is moderate or high.~~

- **CAD Risk—Low**  
10-year absolute coronary or cardiovascular risk less than 10%.
- **CAD Risk—Moderate**  
10-year absolute coronary or cardiovascular risk between 10% and 20%.
- **CAD Risk—High**  
10-year absolute coronary or cardiovascular risk of greater than 20%.

## Websites for Global Cardiovascular Risk Calculators\*

\*Patients who have known CAD are already at high global risk and are not applicable to the calculators (~~Arnet, 2019; D'Agostino, 2008; Goff, 2014; McClelland, 2015; Ridker, 2007~~).<sup>31-35</sup>

Risk Calculator	Link to Online Calculator
Framingham Cardiovascular Risk	<a href="https://reference.medscape.com/calculator/framingham-cardiovascular-disease-risk">https://reference.medscape.com/calculator/framingham-cardiovascular-disease-risk</a>
Reynolds Risk Score Can use if no diabetes Unique for use of family history	<a href="http://www.reynoldsriskscore.org/">http://www.reynoldsriskscore.org/</a>
Pooled Cohort Equation	<a href="http://clincalc.com/Cardiology/ASCVD/PooledCohort.aspx?example">http://clincalc.com/Cardiology/ASCVD/PooledCohort.aspx?example</a>
ACC/AHA Risk Calculator	<a href="http://tools.acc.org/ASCVD-Risk-Estimator/">http://tools.acc.org/ASCVD-Risk-Estimator/</a>
MESA Risk Calculator With addition of Coronary Artery Calcium Score, for CAD-only risk	<a href="https://www.mesa-nhlbi.org/MESACHDRisk/MesaRiskScore/RiskScore.aspx">https://www.mesa-nhlbi.org/MESACHDRisk/MesaRiskScore/RiskScore.aspx</a>

## Definitions of Coronary Artery Disease<sup>2, 23, 27, 36, 37</sup>

(~~Fihn, 2012; Mintz, 2016; Montalescot, 2013; Patel, 2017; Tobis, 2007~~)

- Percentage stenosis refers to the reduction in diameter stenosis when angiography is the method and refers to cross-sectional narrowing when IVUS (intravascular ultrasound) is the method of determination
- Coronary artery calcification is a marker of risk, as measured by Agatston score on coronary artery calcium imaging. ~~It is not a diagnostic tool so much as it is a risk stratification tool.~~ Its incorporation into Global Risk can be achieved by using the MESA risk calculator.
- Ischemia-producing disease (also called hemodynamically or functionally significant disease, for which revascularization might be appropriate), generally implies at least one of the following:
  - Suggested by percentage diameter stenosis  $\geq 70\%$  by angiography; ~~borderline lesions are 40–70%~~<sup>27, 36</sup> (~~Fihn, 2012; Tobis, 2007~~) intermediate lesions are 50 – 69%<sup>38</sup>
  - For a left main artery, suggested by a percentage stenosis  $\geq 50\%$  or minimum lumen cross-sectional area on IVUS  $\leq 6$  square mm<sup>27, 37, 39</sup> (~~Fihn, 2012; Lofti, 2018; Mintz, 2016~~)

- FFR (fractional flow reserve)  $\leq 0.80$  for a major vessel<sup>37, 39</sup> (Lofti, 2018; Mintz, 2016)
- iFR (instantaneous wave free ratio)  $\leq 0.89$  for a major vessel<sup>39-42</sup> (Davies, 2017; Gotberg, 2017; Lofti, 2018)
- A major vessel would be a coronary vessel that would be amenable to revascularization, if indicated. This assessment is made based on the diameter of the vessel and/or the extent of myocardial territory served by the vessel
- FFR (fractional flow reserve) is the distal to proximal pressure ratio across a coronary lesion during maximal hyperemia induced by either intravenous or intracoronary adenosine. Less than or equal to 0.80 is considered a significant reduction in coronary flow
- iFR (instantaneous wave free ratio)  $\leq 0.89$  for a major vessel<sup>40-42</sup> (Davies, 2017; Gotberg, 2017)

### Anginal Equivalent<sup>27, 29, 40</sup>

(Fihn, 2012; Moya, 2009; Shen, 2017)

Development of an anginal equivalent (e.g., shortness of breath, fatigue, or weakness) either with or without prior coronary revascularization should be based upon the documentation of reasons to suspect that symptoms other than chest discomfort are not due to other organ systems (e.g., dyspnea due to lung disease, fatigue due to anemia). This may include respiratory rate, oximetry, lung exam, etc. (as well as d-dimer, chest CT(A), and/or PFTs, when appropriate), and then incorporated into the evaluation of coronary artery disease as would chest discomfort. Syncope per se is not an anginal equivalent.

### Surgical risk delineation

- High risk surgery: Aortic and other major vascular surgery, Peripheral vascular surgery, Anticipated prolonged surgical procedures associated with large fluid shifts and/or blood loss
- Intermediate risk surgery: Carotid endarterectomy, Head and neck surgery, Intraperitoneal and intrathoracic surgery, Orthopedic surgery, Prostate surgery
- Low risk surgery: Endoscopic procedures, superficial procedure, cataract surgery, Breast surgery

## Abbreviations

AAD	Antiarrhythmic drug
ADLs	Activities of daily living
BSA	Body surface area in square meters
<b>CABG</b>	<b>Coronary artery bypass grafting surgery</b>
<b>CAC</b>	<b>Coronary artery calcium</b>
CAD	Coronary artery disease
<b>CCTA</b>	<b>Coronary computed tomography angiography</b>
<b>CMR</b>	<b>Cardiovascular magnetic resonance imaging</b>
<b>CT(A)</b>	<b>Computed tomography (angiography)</b>
<b>DTS</b>	<b>Duke Treadmill Score</b>
ECG	Electrocardiogram
FFR	Fractional flow reserve
<b>HCM</b>	<b>Hypertrophic cardiomyopathy</b>
<b>IVUS</b>	<b>Intravascular ultrasound</b>
LBBB	Left bundle-branch block
LVEF	Left ventricular ejection fraction
LVH	Left ventricular hypertrophy
<b>LVOT</b>	<b>Left ventricular outflow tract</b>
<b>MI</b>	<b>Myocardial infarction</b>
<b>MESA</b>	<b>Multi-Ethnic Study of Atherosclerosis</b>
MET	Estimated metabolic equivalent of exercise
<b>MI</b>	<b>Myocardial infarction</b>
MPI	Myocardial perfusion imaging
<b>MR</b>	<b>Mitral regurgitation</b>
<b>MS</b>	<b>Mitral stenosis</b>
<b>PCI</b>	<b>Percutaneous coronary intervention</b>
<b>PET</b>	<b>Positron emission tomography</b>
PFT	Pulmonary function test
PVCs	Premature ventricular contractions
SE	Stress echocardiography
<b>TTE</b>	<b>Transthoracic echocardiography</b>
VT	Ventricular tachycardia
VF	Ventricular fibrillation
WPW	<b>Wolff-Parkinson-White</b> <del>Wolf Parkinson White</del>

## POLICY HISTORY

Date	Summary
<b><u>February 2022</u></b>	<ul style="list-style-type: none"> <li><del>Brought</del><b>Moved</b> the sentence regarding utilization of suitable alternatives such as Stress Echocardiography and MPI to the <a href="#">General Information section</a></li> </ul>

	<ul style="list-style-type: none"> <li>• <u>Clarified “intermediate lesions are 50-69%” for ischemia-producing disease</u></li> <li>• <u>Placed Link to Overview Section in General Information</u></li> <li>— <del>Added stress imaging approval for calcium score &gt; 100 with low to intermediate probability symptoms</del></li> <li>• <u>Deleted the requirement for diabetes when calcium score &gt; 400 for stress imaging</u></li> <li>• <u>Added Calcium score section:</u> <ul style="list-style-type: none"> <li>○ <u>Added stress imaging approval for calcium score &gt; 100 with symptoms consistent with low to intermediate pretest probability</u></li> </ul> </li> <li>• <u>Changed preoperative guideline to include intermediate risk surgery with one or more risk factors AND documentation of an inability to walk (or &lt;4 METs) AND there has not been an imaging stress test within 1 year</u></li> <li>• <u>Changed solid organ transplant guideline to include stem cell transplant and “any” organ transplant</u> <ul style="list-style-type: none"> <li>— <u>Added definition of surgical risk to preop guidelines</u></li> </ul> </li> <li>— <del>Changed preoperative guideline to An Intermediate or high risk surgery with of one or more risk factors AND documentation of an inability to walk (or &lt;4 METs) AND there has not been an imaging stress test within 1 year*</del></li> <li>— <del>Changed solid organ transplant guideline</del></li> <li>— <del>Added risk factors for preop guidelines and definition of surgical risk to overview section</del></li> <li>• <del></del></li> <li>• <u>In Background section clarified the requirement for description of chest pain by adding sentence “The medical record should provide enough detail to establish the type of chest pain.”</u></li> <li>• <u>Added definition of Q waves</u></li> <li>• <u>Deleted sentence regarding calcium scoring within the Global Risk Section</u></li> <li>• <u>Deleted sentence regarding using calcium score solely for risk stratification</u></li> <li>— <del>Reference to IFR deleted</del> <u>Deleted IFR references</u></li> <li>• <u>Transplant evaluation modified</u></li> </ul>
March 2021	<ul style="list-style-type: none"> <li>• Wording changes to align with MPI</li> <li>• Added annual studies for patients on Flecainide</li> <li>• Added indication for Ca score in diabetic &gt; 40 and calcium score &gt; 400 with reference added</li> <li>• Added indicatios and reference for hypertrophic cardiomyopathy</li> </ul>

	<ul style="list-style-type: none"> <li>Removed BMI &gt; 40 as indication to allow MPI over SE</li> </ul>
August 2020	<ul style="list-style-type: none"> <li>For asymptomatic patients without a history of CAD, the wording for previously unevaluated was changed for Q waves and complete BBB</li> <li>For newly diagnosed systolic heart failure (EF &lt; 50%), when invasive coronary angiography has not been performed, especially when symptoms or signs of ischemia are present or suspected were further defined to state such as: <ul style="list-style-type: none"> <li>Chest pain</li> <li>EKG changes such as new ST segment depression or T wave inversions</li> <li>New wall motion abnormalities</li> </ul> </li> <li>Stress echo with Doppler indication further defined as exertional shortness of breath which suggests the amount of MS is worse than is seen on the resting echocardiogram</li> <li>After the first five years post cardiac transplantation, patients with transplant coronary vasculopathy can be screened annually with ONE of the following: <ul style="list-style-type: none"> <li>MPI</li> <li>SE</li> <li>Left heart catheterization</li> </ul> </li> <li></li> </ul>
March 2020	<ul style="list-style-type: none"> <li>Added general information section as Introduction which outlines requirements for documentation of pertinent office notes by a licensed clinician, and inclusion of laboratory testing and relevant imaging results for case review</li> <li>Added clarification of repeat testing in a patient with new or worsening symptoms and negative result at least one year prior to include the statement “AND meets one of the criteria above”</li> <li>Added clarification of frequent PVCs under ventricular arrhythmias which states defined as greater than or equal to 30/hour to include “on remote monitoring”</li> <li>Edited indication of planning for solid organ transplantation to remove the requirement of limited functional capacity but maintaining requirement of ≥ 3 listed risk factors</li> <li>Added edits to the Coronary Artery Disease definition section</li> <li>Updated and added new references</li> </ul>
November 2019	<ul style="list-style-type: none"> <li>Added CPT code +93356</li> </ul>

July 2019	<ul style="list-style-type: none"> <li>• Stress echo for suspected CAD deleted the following indication: Repeat testing in patient with recurrent symptomatic presentation and negative result over 2 years ago</li> <li>• Added indications: ‘For assessment of hemodynamic significance due to atherosclerosis or following radiation therapy to the anterior or left chest, at 5 years post initiation inception of radiation and every 5 years thereafter’; and ‘Following radiation therapy to the anterior or left chest, at 5 years post initiation inception of radiation and every 5 years thereafter’</li> <li>• Removed secondary mitral regurgitation indication under doppler evaluation section</li> <li>• Clarified indication as follows: Routine follow-up of asymptomatic or stable symptoms when last invasive or non-invasive assessment of coronary disease showed hemodynamically significant CAD (ischemia on stress test or FFR less than or equal to 0.80 or stenosis greater than or equal to 70% of a major vessel) over two years ago without intervening coronary revascularization is an appropriate indication for stress imaging (MPI or SE) in patients if it will alter management</li> </ul>
-----------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

## REFERENCES

~~Acampa W, Assante R, Zampella E. The role of treadmill exercise testing in women. *J Nucl Cardiol*. 2016 Oct; 23(5):991-996.~~

~~Al Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *JACC*. 2018; 72(14):e91-e220.~~

~~Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019; 74:e177-232.~~

~~Baumgartner H, Falk V, Bax JJ, et al. 2017 ESC/EACTS guidelines for the management of valvular heart disease, The Task Force for the Management of Valvular Heart Disease of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). *Euro Heart J*. 2017; 38:2739-2791.~~

~~Bonow RO, O’Gara PT, Adams DH, et al. 2020 Focused Update of the 2017 ACC Expert Consensus Decision Pathway on the Management of Mitral Regurgitation. *J Am Coll Cardiol*. Epub 2020 Feb 14.~~

~~Budoff MJ, Raggi P, Beller GA, et al. Noninvasive Cardiovascular Risk Assessment of the Asymptomatic Diabetic Patient: The Imaging Council of the American College of Cardiology. *JACC Cardiovasc Imaging*. 2016 Feb; 9(2):176-92.~~

~~Cha Y, Lee GK, Klarich KW, et al. Advances in Arrhythmia and Electrophysiology. Premature Ventricular Contraction-Induced Cardiomyopathy, A Treatable Condition. *Circulation: Arrhythmia and Electrophysiology*. 2012; 5:229-236. Available at: <http://circep.ahajournals.org/content/5/1/229.full>~~

~~Crea F, Camici PG, Merz CN. Coronary microvascular dysfunction: an update. *Euro Heart J*. 2013; 35(17):1101-1111. Available at: <http://eurheartj.oxfordjournals.org/content/early/2013/12/21/eurheartj.eht513>~~

~~D’Agostino RB Sr, Vasan RS, Pencina MJ, et al. General Cardiovascular Risk Profile for Use in Primary Care: The Framingham Heart Study. *Circulation*. 2008; 117:743-753.~~

~~Doherty JU, Kort S, Mehran R, et al. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2017 Appropriate Use Criteria for Multimodality Imaging in Valvular Heart Disease. A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American~~



Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *JACC*. 2017; 70(13):1647-1672.

Doherty JU, Kort S, Mehran R, et al. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2019 Appropriate Use Criteria for Multimodality Imaging in the Assessment of Cardiac Structure and Function in Nonvalvular Heart Disease : A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and the Society of Thoracic Surgeons. *J Nucl Cardiol*. 2019; 26(4):1392-1413.

Douglas PS et al. ACCF/ASE/AHA/ASNC/HFSA/HRS/SCAI/SCCM/SCCT/SCMR 2011 Appropriate Use Criteria for Echocardiography. *J Am Soc Echocardiography*. 2011; 24:229-67.

Einstein, A. Effects of radiation exposure from cardiac imaging: how good are the data? *Journal of the American College of Cardiology*. 2012; 59(6):553-565. Available at: <http://content.onlinejacc.org/cgi/content/short/59/6/553>

Fazel R, Dilsizian V, Einstein, AJ, et al. ASNC INFORMATION STATEMENT, Strategies for defining an optimal risk-benefit ratio for stress myocardial perfusion SPECT. *Journal of Nuclear Cardiology*. Published Online: 24 March 2011:1-8. Available at: <https://www.asnc.org/files/Optimal%20Risk-Benefit%20Ratio%20for%20SPECT.pdf>

Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the diagnosis and management of patients with stable ischemic heart disease: A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *Circulation*. 2012; 126(25): e354-471.

Fleischer LA, Fleischmann KE, Auerbach AD, et al. ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery. A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014; 64(22): e77-e137. Available at: <http://content.onlinejacc.org/article.aspx?articleid=1893784>

Fleischmann KE, Hunink MG, Kuntz KM, et al. Exercise echocardiography or exercise SPECT imaging? A meta-analysis of diagnostic test performance. *JAMA*. 1998; 280:913-20.

Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients with Lower Extremity Peripheral Artery Disease: A Report of the American College of

Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *JACC*. Nov 8, in press. Available at: [www.onlinejacc.org/content/early/2016/11/23/j.jacc.2016.11.007](http://www.onlinejacc.org/content/early/2016/11/23/j.jacc.2016.11.007)

Goff DC, Lloyd Jones, DM, Bennett G, et al. 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk, A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines Endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation, American Society for Preventive Cardiology, American Society of Hypertension, Association of Black Cardiologists, National Lipid Association, Preventive Cardiovascular Nurses Association, and Women Heart: The National Coalition for Women With Heart Disease. *JACC*. 2014; 63(25): 2935-2959.

Grani C, Buechel RR, Kaufmann PA, et al. Multimodality Imaging in Individuals with Anomalous Coronary Arteries. *JACC*. 2017; 10(4): 471-581.

Heijenbroek-Kal MH, Fleischmann KE, Hunink MG. Stress echocardiography, stress single-photon-emission-computed tomography and electron beam-computed tomography for the assessment of coronary artery disease: a meta-analysis of diagnostic performance. *Am Heart J*. 2007 Sep; 154(3):415-23.

Henzlova MJ, Duvall WL, Einstein, AJ, et al, ASNC imaging guidelines for SPECT nuclear cardiology procedures: Stress, protocols, and tracers, *J Nucl Cardiol*. 2016; 23:606-39. Available at: <https://link.springer.com/content/pdf/10.1007/s12350-015-0387-x.pdf>

Hirshfeld JW, Ferrari VA, Bengel FM, et al. 2018 ACC/HRS/NASCI/SCAI/SCCT Expert Consensus Document on Optimal Use of Ionizing Radiation in Cardiovascular Imaging: Best Practices for Safety and Effectiveness A Report of the American College of Cardiology Task Force on Expert Consensus Decision Pathways Developed in Collaboration with Mended Hearts. *JACC*. in press, May, 2018: e1-e69. Available at: <http://www.onlinejacc.org/content/early/2018/04/30/j.jacc.2018.02.016>

Hussain MA, Al-Omram M, Creager MA, et al. Antithrombotic Therapy for Peripheral Artery Disease, Recent Advances. *JACC*. 2018; 71(21):2450-2467.

Lancellotti P, Knemo VT, Badano LP, et al. Expert consensus for multi-modality imaging evaluation of cardiovascular complications of radiotherapy in adults: a report from the European Association of Cardiovascular Imaging and the American Society of Echocardiography. *Euro Heart J – Cardiovas Imaging*. 2013; 14:721-740.

Lentine KL, Costa SP, Weir MR. Cardiac disease evaluation and management among kidney and liver transplantation candidates. *JACC*. 2012; 60(5): 434-480.

Lotfi A, Davies JE, Fearon WF, et al. Focused update of expert consensus statement: Use of invasive assessments of coronary physiology and structure: A position statement of the society of cardiac angiography and interventions. *Catheter Cardiovasc Interv*. 2018 Jul 3.

Mark DB, Hlatky MA, Harrell FE Jr, et al. Exercise treadmill score for predicting prognosis in coronary artery disease. *Ann Intern Med.* 1987; 106(6):793-800.

Marwick, TH. Stress echocardiography. *Heart.* 2003; 89(1): 113-118. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1767520/>.

McClelland RL, Jorgensen NW, Budoff M, et al. 10-year coronary heart disease risk prediction using coronary artery calcium and traditional risk factors: Derivation in the MESA (Multi-Ethnic Study of Atherosclerosis) with validation in the HNR (Heinz Nixdorf Recall) Study and the DHS (Dallas Heart Study). *JACC.* 2015; 66(15):1643-53.

McCrindle BW, Rowley AH, Newburger JW, et al. Diagnosis, treatment, and long-term management of Kawasaki disease: A scientific statement for health professionals from the American Heart Association. *Circulation.* 2017; 135(17): e927.

Metz, LD, Beattie, M, Hom, R, et al. The prognostic value of normal exercise myocardial perfusion imaging and exercise echocardiography: A meta-analysis. *JACC.* 2007; 49:227-237. Available at: <http://content.onlinejacc.org/cgi/reprint/49/2/227.pdf>

Miller TD, Askew JW, Anavekar NS. Noninvasive Stress Testing for Coronary Artery Disease. *Heart Fail Clin.* 2016; 12(1):65-82.

Mintz GS. IVUS in PCI Guidance. *Am Coll Cardiol*, (online website) June 13, 2016. Available at: <http://www.acc.org/latest-in-cardiology/articles/2016/06/13/10/01/ivus-in-pci-guidance>.

Montalescot G, Sechtem U, Achenbach S, et al. 2013 ESC guidelines on the management of stable coronary artery disease: The Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *European Heart Journal.* 2013; 34(38): 2949-3003. Available at: <https://academic.oup.com/eurheartj/article/34/38/2949/442952>.

Moya A, Sutton R, Ammirati F, et al. Guidelines for the diagnosis and management of syncope: Task Force for the Diagnosis and Management of Syncope. *Euro Heart J.* 2009; 30: 2631-71. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3295536/>.

Mukundan S, Travin MI, Levsky JM, et al. Does ischemic burden on stress testing influence patient survival in subjects with known severe multi-vessel CAD? *Am J Cardiovasc Dis.* 2017;7(2):48-52.

Nishimura RA, Otto CM, Bonow RO, et al. 2014 AHA/ACC Guideline for the Management of Patients with Valvular Heart Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2014; 63(22):e57-e185.

~~Ommen ST, Mital S, Burke MA, et al. 2020 AHA/ACC Guideline for the Diagnosis and Treatment of Patients With Hypertrophic Cardiomyopathy: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2020 Dec 22; 76(25):3022-55.~~

~~Patel AY, Eagle KA, Vaishnava P. Cardiac Risk of Noncardiac Surgery. *J Am Coll Cardiol*. 2015; 66(19):2140-2148. Available at: <http://content.onlinejacc.org/article.aspx?articleid=2468532>.~~

~~Patel MR, Bailey SR, Bonow RO, et al. ACCF/SCAI/AATS/AHA/ASE/ASNC/HFSA/HRS/SCCM/SCCT/SCMR/STS 2012 Appropriate Use Criteria for Diagnostic Catheterization: A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *JACC*. 2012; 59(22):1995-2027.~~

~~Patel MR, Calhoon JH, Dehmer GJ, et al. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2017 Appropriate Use Criteria for Coronary Revascularization in Patients with Stable Ischemic Heart Disease. *JACC*. 2017; 69(17):2212-2241.~~

~~Patel MR, White RD, Abbara S, et al. 2013 ACCF/ACR/ASE/ASNC/SCCT/SCMR Appropriate Utilization of Cardiovascular Imaging in Heart Failure, A Joint Report of the American College of Radiology Appropriateness Criteria Committee and the American College of Cardiology Foundation Appropriate Use Criteria Task Force. *JACC*. 2013; 61(21): 2207-2231.~~

~~Pellikka PA, Nagueh SF, Elhendy AA, et al. American Society of Echocardiography Recommendations for Performance, Interpretation, and Application of Stress Echocardiography. *Journal of the American Society of Echocardiography*. 2007; 20(9):1021-41.~~

~~Reiffel JA, Camm AJ, Belardinelli L, et al. The HARMONY trial: Combined ranolazine and dronedarone in the management of paroxysmal atrial fibrillation: Mechanistic and therapeutic synergism. *Circ Arrhythm Electrophysiol*. 2015; 8(5):1048.~~

~~Ridker PM, Buring JE, Rifai N, et al. NIH Estimate of 10 Year coronary artery disease risk from Framingham Risk Score: Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds Risk Score. *JAMA*. 2007; 297(6):611-619. Available at: <http://jama.jamanetwork.com/article.aspx?articleid=205528>.~~

~~Schinkel AFL, Bax JJ, Geleijnse ML, et al. Noninvasive evaluation of ischaemic heart disease: myocardial perfusion imaging or stress echocardiography? *Euro Heart J*. 2003; 24(9):789-800. Available at: [https://doi.org/10.1016/S0195-668X\(02\)00634-6](https://doi.org/10.1016/S0195-668X(02)00634-6).~~

~~Scott-Moncrieff, A, Yang J, Levine D, et al. Real-world estimated effective radiation doses from commonly used cardiac testing and procedural modalities. *The Canadian Journal of Cardiology*. 2011; 27(5):613-618. Available at:~~

~~[http://www.unboundmedicine.com/medline/ebm/record/21652170/abstract/Real\\_world\\_estimated\\_effective\\_radiation\\_doses\\_from\\_commonly\\_used\\_cardiac\\_testing\\_and\\_procedural\\_modalities](http://www.unboundmedicine.com/medline/ebm/record/21652170/abstract/Real_world_estimated_effective_radiation_doses_from_commonly_used_cardiac_testing_and_procedural_modalities).~~

~~Shen W, Sheldon RS, Benditt DG, et al. 2017 ACC/AHA/HRS Guideline for the Evaluation and Management of Patients with Syncope: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *JACC*. 2017; 70(5):620-663.~~

~~Sicari R, Cortigiani L. The clinical use of stress echocardiography in ischemic heart disease. *Cardiovascular Ultrasound*. 2017; 15(7):1-16. Available at:~~

~~<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5361820/>.~~

~~Sicari, R. Stress echocardiography expert consensus statement, European Association of Echocardiography (EAE) (a registered branch of the ESC). *European Journal of Echocardiography*. 2008; 9:415-443. Available at:~~

~~[https://www.escardio.org/static\\_file/Escardio/Subspecialty/EACVI/position\\_papers/eae-sicari-stress-echo.pdf](https://www.escardio.org/static_file/Escardio/Subspecialty/EACVI/position_papers/eae-sicari-stress-echo.pdf).~~

~~Steiner J, Rodes-Cabau J, Holmes D, et al. Mechanical intervention for aortic valve stenosis in patients with heart failure and reduced ejection fraction. *JACC*. 2017; 70(24):3026-3041. Available at: <http://www.onlinejacc.org/content/70/24/3026>.~~

~~Tang K, Wang L, Shi R, et al. The role of myocardial perfusion imaging in evaluating patients with myocardial bridging. *J Nucl Cardiol*. 2011; 18(1):117.~~

~~Tobis J, Azarbal B, Slavin L. Assessment of Intermediate Severity Coronary Lesions in the Catheterization Laboratory. *JACC*. 2007; 49(8):839-848.~~

~~Wolk MJ, Bailey SR, Doherty JU, et al. ACCF/AHA/ASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 Multimodality Appropriate Use Criteria for the Detection and Risk Assessment of Stable Ischemic Heart Disease A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *Journal of the American College of Cardiology*. 2014;63(4): 380-406. Available at:~~

~~<http://content.onlinejacc.org/article.aspx?articleid=1789799>.~~

~~Yancy C, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA Guideline for the Management of Heart~~

~~Failure, A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *JACC*. 2013; 62(16): e147-237.~~

~~Yao S, Qureshi E, Sherrid M, et al. Practical Application in Stress Echocardiography, Risk Stratification and Prognosis in Patients with Known or suspected Ischemic Heart disease. *JACC*. 2003; 42(6):1084-90.~~

~~Zhang Y, Li X, Segars WP, et al. Comparison of patient specific dose metrics between chest radiography, tomosynthesis, and CT for adult patients of wide-ranging body habitus. *Med. Phys.* 2014; 41(2): 1-12. Available at: [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3985882/pdf/MPHYA6-000041-023901\\_1.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3985882/pdf/MPHYA6-000041-023901_1.pdf).~~

~~Zimetbaum PJ. Nonsustained ventricular tachycardia: Clinical manifestation, evaluation, and management, UpToDate. Waltham, MA; April, 2018. Available at: [https://www.uptodate.com/contents/nonsustained-ventricular-tachycardia-clinical-manifestations-evaluation-and-management?search=nonsustained%20ventricular%20tachycardia&source=search\\_result&selectedTitle=1~94&usage\\_type=default&display\\_rank=1#H16](https://www.uptodate.com/contents/nonsustained-ventricular-tachycardia-clinical-manifestations-evaluation-and-management?search=nonsustained%20ventricular%20tachycardia&source=search_result&selectedTitle=1~94&usage_type=default&display_rank=1#H16) Retrieved May 18, 2018~~

**~~Reviewed / Approved by NIA Clinical Guideline Committee~~**

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

1. Gulati M, Levy PD, Mukherjee D, et al. 2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Cardiovasc Comput Tomogr*. Dec 01 2021;doi:10.1016/j.jcct.2021.11.009
2. Patel MR, Calhoon JH, Dehmer GJ, et al. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2017 Appropriate Use Criteria for Coronary Revascularization in Patients With Stable Ischemic Heart Disease: A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. May 2 2017;69(17):2212-2241. doi:10.1016/j.jacc.2017.02.001
3. Wolk MJ, Bailey SR, Doherty JU, et al. ACCF/AHA/ASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 multimodality appropriate use criteria for the detection and risk assessment of stable ischemic heart disease: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Feb 4 2014;63(4):380-406. doi:10.1016/j.jacc.2013.11.009
4. Budoff MJ, Raggi P, Beller GA, et al. Noninvasive Cardiovascular Risk Assessment of the Asymptomatic Diabetic Patient: The Imaging Council of the American College of Cardiology. *JACC Cardiovasc Imaging*. Feb 2016;9(2):176-92. doi:10.1016/j.jcmg.2015.11.011
5. Aguilar-Salinas CA, Rojas R, Gomez-Perez FJ, et al. Analysis of the agreement between the World Health Organization criteria and the National Cholesterol Education Program-III definition of the metabolic syndrome: results from a population-based survey. *Diabetes Care*. May 2003;26(5):1635.
6. Doherty JU, Kort S, Mehran R, Schoenhagen P, Soman P. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2019 Appropriate Use Criteria for



- Multimodality Imaging in the Assessment of Cardiac Structure and Function in Nonvalvular Heart Disease : A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and the Society of Thoracic Surgeons. *J Nucl Cardiol*. Aug 2019;26(4):1392-1413. doi:10.1007/s12350-019-01751-7
7. Doherty JU, Kort S, Mehran R, et al. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2019 Appropriate Use Criteria for Multimodality Imaging in the Assessment of Cardiac Structure and Function in Nonvalvular Heart Disease : A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and the Society of Thoracic Surgeons. *J Nucl Cardiol*. Aug 2019;26(4):1392-1413. doi:10.1007/s12350-019-01751-7
8. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS Guideline for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. Oct 2 2018;72(14):e91-e220. doi:10.1016/j.jacc.2017.10.054
9. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS Guideline for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation*. 09 25 2018;138(13):e272-e391. doi:10.1161/CIR.0000000000000549
10. Reiffel JA, Camm AJ, Belardinelli L, et al. The HARMONY Trial: Combined Ranolazine and Dronedarone in the Management of Paroxysmal Atrial Fibrillation: Mechanistic and Therapeutic Synergism. *Circ Arrhythm Electrophysiol*. Oct 2015;8(5):1048-56. doi:10.1161/circep.115.002856
11. Gräni C, Buechel RR, Kaufmann PA, Kwong RY. Multimodality Imaging in Individuals With Anomalous Coronary Arteries. *JACC Cardiovasc Imaging*. Apr 2017;10(4):471-481. doi:10.1016/j.jcmg.2017.02.004
12. Tang K, Wang L, Shi R, et al. The role of myocardial perfusion imaging in evaluating patients with myocardial bridging. *J Nucl Cardiol*. Feb 2011;18(1):117-22. doi:10.1007/s12350-010-9303-6
13. McCrindle BW, Rowley AH, Newburger JW, et al. Diagnosis, Treatment, and Long-Term Management of Kawasaki Disease: A Scientific Statement for Health Professionals From the American Heart Association. *Circulation*. Apr 25 2017;135(17):e927-e999. doi:10.1161/cir.0000000000000484
14. Lancellotti P, Nkomo VT, Badano LP, et al. Expert consensus for multi-modality imaging evaluation of cardiovascular complications of radiotherapy in adults: a report from the European Association of Cardiovascular Imaging and the American Society of Echocardiography. *Eur Heart J Cardiovasc Imaging*. Aug 2013;14(8):721-40. doi:10.1093/ehjci/jet123



15. Bonow RO, O'Gara PT, Adams DH, et al. 2020 Focused Update of the 2017 ACC Expert Consensus Decision Pathway on the Management of Mitral Regurgitation: A Report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol*. May 5 2020;75(17):2236-2270. doi:10.1016/j.jacc.2020.02.005
16. Nishimura RA, Otto CM, Bonow RO, et al. 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Jun 10 2014;63(22):e57-185. doi:10.1016/j.jacc.2014.02.536
17. Baumgartner H, Falk V, Bax JJ, et al. 2017 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J*. Sep 21 2017;38(36):2739-2791. doi:10.1093/eurheartj/ehx391
18. Steiner J, Rodés-Cabau J, Holmes DR, Jr., LeWinter MM, Dauerman HL. Mechanical Intervention for Aortic Valve Stenosis in Patients With Heart Failure and Reduced Ejection Fraction. *J Am Coll Cardiol*. Dec 19 2017;70(24):3026-3041. doi:10.1016/j.jacc.2017.10.040
19. Ommen SR, Mital S, Burke MA, et al. 2020 AHA/ACC Guideline for the Diagnosis and Treatment of Patients With Hypertrophic Cardiomyopathy: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. Dec 22 2020;76(25):3022-3055. doi:10.1016/j.jacc.2020.08.044
20. Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*. Dec 9 2014;64(22):e77-137. doi:10.1016/j.jacc.2014.07.944
21. Patel AY, Eagle KA, Vaishnava P. Cardiac risk of noncardiac surgery. *J Am Coll Cardiol*. Nov 10 2015;66(19):2140-2148. doi:10.1016/j.jacc.2015.09.026
22. Kristensen SD, Knuuti J, Saraste A, et al. 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *Eur Heart J*. Sep 14 2014;35(35):2383-431. doi:10.1093/eurheartj/ehu282
23. Montalescot G, Sechtem U, Achenbach S, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. Oct 2013;34(38):2949-3003. doi:10.1093/eurheartj/ehs296
24. Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*. Dec 09 2014;64(22):e77-137. doi:10.1016/j.jacc.2014.07.944
25. Velasco A, Reyes E, Hage FG. Guidelines in review: Comparison of the 2014 ACC/AHA guidelines on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery and the 2014 ESC/ESA guidelines on noncardiac surgery: Cardiovascular assessment and management. *J Nucl Cardiol*. 02 2017;24(1):165-170. doi:10.1007/s12350-016-0643-8

26. Lentine KL, Costa SP, Weir MR, et al. Cardiac disease evaluation and management among kidney and liver transplantation candidates: a scientific statement from the American Heart Association and the American College of Cardiology Foundation. *J Am Coll Cardiol*. Jul 31 2012;60(5):434-80. doi:10.1016/j.jacc.2012.05.008
27. Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *Circulation*. Dec 18 2012;126(25):e354-471. doi:10.1161/CIR.0b013e318277d6a0
28. Henzlova MJ, Duvall WL, Einstein AJ, Travin MI, Verberne HJ. ASNC imaging guidelines for SPECT nuclear cardiology procedures: Stress, protocols, and tracers. *J Nucl Cardiol*. Jun 2016;23(3):606-39. doi:10.1007/s12350-015-0387-x
29. Shen WK, Sheldon RS, Benditt DG, et al. 2017 ACC/AHA/HRS Guideline for the Evaluation and Management of Patients With Syncope: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. Aug 1 2017;70(5):620-663. doi:10.1016/j.jacc.2017.03.002
30. Mark DB, Hlatky MA, Harrell FE, Jr., Lee KL, Califf RM, Pryor DB. Exercise treadmill score for predicting prognosis in coronary artery disease. *Ann Intern Med*. Jun 1987;106(6):793-800. doi:10.7326/0003-4819-106-6-793
31. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. Sep 10 2019;74(10):e177-e232. doi:10.1016/j.jacc.2019.03.010
32. D'Agostino RB, Sr., Vasan RS, Pencina MJ, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation*. Feb 12 2008;117(6):743-53. doi:10.1161/circulationaha.107.699579
33. Goff DC, Jr., Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Jul 1 2014;63(25 Pt B):2935-2959. doi:10.1016/j.jacc.2013.11.005
34. McClelland RL, Jorgensen NW, Budoff M, et al. 10-Year Coronary Heart Disease Risk Prediction Using Coronary Artery Calcium and Traditional Risk Factors: Derivation in the MESA (Multi-Ethnic Study of Atherosclerosis) With Validation in the HNR (Heinz Nixdorf Recall) Study and the DHS (Dallas Heart Study). *J Am Coll Cardiol*. Oct 13 2015;66(15):1643-53. doi:10.1016/j.jacc.2015.08.035
35. Ridker PM, Buring JE, Rifai N, Cook NR. Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds Risk Score. *Jama*. Feb 14 2007;297(6):611-9. doi:10.1001/jama.297.6.611
36. Tobis J, Azarbal B, Slavin L. Assessment of intermediate severity coronary lesions in the catheterization laboratory. *J Am Coll Cardiol*. Feb 27 2007;49(8):839-48. doi:10.1016/j.jacc.2006.10.055

37. Mintz G. IVUS in PCI Guidance. American College of Cardiology. Updated June 13, 2016. Accessed October 22, 2021. <https://www.acc.org/latest-in-cardiology/articles/2016/06/13/10/01/ivus-in-pci-guidance>
38. Patel MR, Bailey SR, Bonow RO, et al. ACCF/SCAI/AATS/AHA/ASE/ASNC/HFSA/HRS/SCCM/SCCT/SCMR/STS 2012 appropriate use criteria for diagnostic catheterization: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, Society of Thoracic Surgeons. *J Thorac Cardiovasc Surg*. Jul 2012;144(1):39-71. doi:10.1016/j.jtcvs.2012.04.013
39. Lotfi A, Davies JE, Fearon WF, Grines CL, Kern MJ, Klein LW. Focused update of expert consensus statement: Use of invasive assessments of coronary physiology and structure: A position statement of the society of cardiac angiography and interventions. *Catheter Cardiovasc Interv*. Aug 1 2018;92(2):336-347. doi:10.1002/ccd.27672
40. Moya A, Sutton R, Ammirati F, et al. Guidelines for the diagnosis and management of syncope (version 2009). *Eur Heart J*. Nov 2009;30(21):2631-71. doi:10.1093/eurheartj/ehp298

## **ADDITIONAL RESOURCES**

1. Acampa W, Assante R, Zampella E. The role of treadmill exercise testing in women. *J Nucl Cardiol*. Oct 2016;23(5):991-996. doi:10.1007/s12350-016-0596-y
2. Lee GK, Klarich KW, Grogan M, Cha YM. Premature ventricular contraction-induced cardiomyopathy: a treatable condition. *Circ Arrhythm Electrophysiol*. Feb 2012;5(1):229-36. doi:10.1161/circep.111.963348
3. Crea F, Camici PG, Bairey Merz CN. Coronary microvascular dysfunction: an update. *Eur Heart J*. May 2014;35(17):1101-11. doi:10.1093/eurheartj/ehf513
4. Doherty JU, Kort S, Mehran R, Schoenhagen P, Soman P. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2017 Appropriate Use Criteria for Multimodality Imaging in Valvular Heart Disease: A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Sep 26 2017;70(13):1647-1672. doi:10.1016/j.jacc.2017.07.732
5. Douglas PS, Garcia MJ, Haines DE, et al. ACCF/ASE/AHA/ASNC/HFSA/HRS/SCAI/SCCM/SCCT/SCMR 2011 Appropriate Use Criteria for Echocardiography. A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Society of Echocardiography, American Heart Association, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm

Society, Society for Cardiovascular Angiography and Interventions, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance American College of Chest Physicians. *J Am Soc Echocardiogr.* Mar 2011;24(3):229-67. doi:10.1016/j.echo.2010.12.008

6. Einstein AJ. Effects of radiation exposure from cardiac imaging: how good are the data? *J Am Coll Cardiol.* Feb 7 2012;59(6):553-65. doi:10.1016/j.jacc.2011.08.079

7. Fazel R, Dilsizian V, Einstein AJ, Ficaro EP, Henzlova M, Shaw LJ. Strategies for defining an optimal risk-benefit ratio for stress myocardial perfusion SPECT. *J Nucl Cardiol.* May 2011;18(3):385-92. doi:10.1007/s12350-011-9353-4

8. Fleischmann KE, Hunink MG, Kuntz KM, Douglas PS. Exercise echocardiography or exercise SPECT imaging? A meta-analysis of diagnostic test performance. *Jama.* Sep 9 1998;280(10):913-20. doi:10.1001/jama.280.10.913

9. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* Mar 21 2017;69(11):e71-e126. doi:10.1016/j.jacc.2016.11.007

10. Heijenbroek-Kal MH, Fleischmann KE, Hunink MG. Stress echocardiography, stress single-photon-emission computed tomography and electron beam computed tomography for the assessment of coronary artery disease: a meta-analysis of diagnostic performance. *Am Heart J.* Sep 2007;154(3):415-23. doi:10.1016/j.ahj.2007.04.061

11. Hirshfeld JW, Jr., Ferrari VA, Bengel FM, et al. 2018 ACC/HRS/NASCI/SCAI/SCCT Expert Consensus Document on Optimal Use of Ionizing Radiation in Cardiovascular Imaging-Best Practices for Safety and Effectiveness, Part 1: Radiation Physics and Radiation Biology: A Report of the American College of Cardiology Task Force on Expert Consensus Decision Pathways Developed in Collaboration With Mended Hearts. *Catheter Cardiovasc Interv.* Aug 1 2018;92(2):203-221. doi:10.1002/ccd.27660

12. Hussain MA, Al-Omran M, Creager MA, Anand SS, Verma S, Bhatt DL. Antithrombotic Therapy for Peripheral Artery Disease: Recent Advances. *J Am Coll Cardiol.* May 29 2018;71(21):2450-2467. doi:10.1016/j.jacc.2018.03.483

13. Marwick TH. Stress echocardiography. *Heart.* 2003;89(1):113-118. doi:10.1136/heart.89.1.113

14. Metz LD, Beattie M, Hom R, Redberg RF, Grady D, Fleischmann KE. The prognostic value of normal exercise myocardial perfusion imaging and exercise echocardiography: a meta-analysis. *J Am Coll Cardiol.* Jan 16 2007;49(2):227-37. doi:10.1016/j.jacc.2006.08.048

15. Miller TD, Askew JW, Anavekar NS. Noninvasive Stress Testing for Coronary Artery Disease. *Heart Fail Clin.* Jan 2016;12(1):65-82. doi:10.1016/j.hfc.2015.08.006

16. Mukundan S, Travin MI, Levsky JM, Liakos MP, Spevack DM. Does ischemic burden on stress testing influence patient survival in subjects with known severe multi-vessel CAD? *Am J Cardiovasc Dis.* 2017;7(2):48-52.

17. Patel MR, Bailey SR, Bonow RO, et al. ACCF/SCAI/AATS/AHA/ASE/ASNC/HFSA/HRS/SCCM/SCCT/SCMR/STS 2012 appropriate use criteria for diagnostic catheterization: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American Heart Association,

- American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. May 29 2012;59(22):1995-2027. doi:10.1016/j.jacc.2012.03.003
18. Patel MR, White RD, Abbara S, et al. 2013 ACCF/ACR/ASE/ASNC/SCCT/SCMR appropriate utilization of cardiovascular imaging in heart failure: a joint report of the American College of Radiology Appropriateness Criteria Committee and the American College of Cardiology Foundation Appropriate Use Criteria Task Force. *J Am Coll Cardiol*. May 28 2013;61(21):2207-31. doi:10.1016/j.jacc.2013.02.005
19. Pellikka PA, Nagueh SF, Elhendy AA, Kuehl CA, Sawada SG. American Society of Echocardiography recommendations for performance, interpretation, and application of stress echocardiography. *J Am Soc Echocardiogr*. Sep 2007;20(9):1021-41. doi:10.1016/j.echo.2007.07.003
20. Schinkel AF, Bax JJ, Geleijnse ML, et al. Noninvasive evaluation of ischaemic heart disease: myocardial perfusion imaging or stress echocardiography? *Eur Heart J*. May 2003;24(9):789-800. doi:10.1016/s0195-668x(02)00634-6
21. Scott-Moncrieff A, Yang J, Levine D, et al. Real-world estimated effective radiation doses from commonly used cardiac testing and procedural modalities. *Can J Cardiol*. Sep-Oct 2011;27(5):613-8. doi:10.1016/j.cjca.2011.01.011
22. Sicari R, Cortigiani L. The clinical use of stress echocardiography in ischemic heart disease. *Cardiovasc Ultrasound*. Mar 21 2017;15(1):7. doi:10.1186/s12947-017-0099-2
23. Sicari R, Nihoyannopoulos P, Evangelista A, et al. Stress echocardiography expert consensus statement: European Association of Echocardiography (EAE) (a registered branch of the ESC). *Eur J Echocardiogr*. Jul 2008;9(4):415-37. doi:10.1093/ejechocard/jen175
24. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Oct 15 2013;62(16):e147-239. doi:10.1016/j.jacc.2013.05.019
25. Yao SS, Qureshi E, Sherriid MV, Chaudhry FA. Practical applications in stress echocardiography: risk stratification and prognosis in patients with known or suspected ischemic heart disease. *J Am Coll Cardiol*. Sep 17 2003;42(6):1084-90. doi:10.1016/s0735-1097(03)00923-9
26. Zhang Y, Li X, Segars WP, Samei E. Comparison of patient specific dose metrics between chest radiography, tomosynthesis, and CT for adult patients of wide ranging body habitus. *Med Phys*. 2014;41(2):023901-023901. doi:10.1118/1.4859315

Reviewed / Approved by NIA Clinical Guideline Committee

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.